

# Doravirine-lamivudine-tenofovir DF (Delstrigo®)

100-300-300 mg – yellow, oval tablet

## What is Delstrigo®?

The Delstrigo® tablet contains 3 different medicines: doravirine, lamivudine, and tenofovir DF. Delstrigo® is used to lower the amount of HIV virus in your body and make your immune system stronger.

## How do I take it?

- Take 1 tablet once daily, at the same time every day, with or without food.

Store in a dry place at room temperature, and keep out of reach of children.

## What if I forget to take it?

It is very important to take Delstrigo® every day to make sure that it keeps working. If you miss a dose, there is a higher chance that the virus will change to a form that is harder to treat. This is called "resistance".

If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, do not take 2 doses at the same time. If you are not sure if you should take your dose, call the pharmacist.

If you stop taking HIV medicine, the amount of HIV virus in your blood will increase.

If you have Hepatitis B, stopping Delstrigo® can make the Hepatitis B worsen.

**Do not stop any of your HIV medicines without talking to your doctor.**

## What should I expect?

Delstrigo® is usually well tolerated. Side effects may include:

- |                       |             |
|-----------------------|-------------|
| • nausea and vomiting | • diarrhea  |
| • trouble sleeping    | • dizziness |
| • fatigue             | • headache  |
| • stomach pain        |             |

Rare, serious side effects may include:

- Allergic reaction: Signs include rash, blisters on skin or mouth, swelling of the face or throat, difficulty breathing, fever, flu-like symptoms
- New or worse kidney problems: Signs include being unable to pass urine or change in amount of urine, blood in urine, weight gain
- Liver problems: Signs include yellow skin or eyes, dark urine, pale stools, or pain on right side below ribs
- Loss of bone strength
- A condition called lactic acidosis: Signs include unusual muscle pain, stomach pain with nausea and vomiting, or a fast heartbeat or fast breathing
- Inflamed pancreas: Symptoms include nausea, vomiting and severe stomach pains

**Always tell your doctor or pharmacist if side effects are bothering you.  
IMMEDIATELY contact your doctor if you have any serious side effects**

It is important to have regular blood tests to watch for side effects and make sure your HIV medicine is working to control the virus.



BRITISH COLUMBIA  
CENTRE for EXCELLENCE  
in HIV/AIDS



Talk to your doctor if you are pregnant or are thinking about becoming pregnant, or if you plan to breastfeed.

## Can I take Delstrigo® with other medicines?

Some medicines interact with Delstrigo®. Drug interactions can stop your treatment from working or may cause unwanted side effects.

Always tell your pharmacist and doctor if you are using any prescription or non-prescription medicines, vitamins, supplements, natural medicines or recreational drugs.

Medicines that may interact with Delstrigo® include:

- Adefovir (Hepsera®) for hepatitis B
- Some anti-inflammatory pain medicines (“NSAIDS”) such as ibuprofen (Advil®) or naproxen (Aleve®) may increase the risk of kidney problems. Low dose, short-term use is usually possible. Talk to your doctor before taking these medicines.
- Some medicines to treat seizures such as phenytoin (Dilantin®)
- Some antibiotics such as rifampin and rifabutin
- St. John's Wort

This is not a complete list of medicines that interact with Delstrigo®.

Always check with your pharmacist before starting any new medicines to make sure they are safe for you.

## What if I have questions?

If you have questions or problems with your medicine, call the St. Paul’s Hospital Ambulatory Pharmacy:

**1-888-511-6222**

(Monday to Friday daytime and urgent issues after hours)

To book an appointment to see a St. Paul’s Hospital pharmacist, call reception:

**604-806-8060**

(Monday to Friday daytime)

This pamphlet is for general information purposes only. It does not provide individual medical or treatment advice and is not a substitute for medical or professional care.