



BRITISH COLUMBIA
CENTRE *for* EXCELLENCE
in HIV/AIDS

HIV MONITORING QUARTERLY REPORT FOR BRITISH COLUMBIA

THIRD QUARTER 2013

UPDATED VERSION: NOV 28, 2014 *

** See foreword*



BC Centre for Disease Control
An agency of the Provincial Health Services Authority



Foreword

As part of the BC Centre for Excellence (BC-CFE) in HIV/AIDS's mandate to evaluate the outcomes of STOP HIV/AIDS programming in BC, we have developed quarterly HIV/AIDS monitoring reports. These reports provide up-to-date data on a variety of key HIV-related surveillance and treatment indicators. Selection of these indicators was achieved through a collaborative process with various Health Authority (HA) representatives. There are six reports in total, one for each HA and one for the province of BC as a whole. In addition, there is a technical report which explains how each HIV indicator is calculated. Data used in these reports come from the British Columbia Centre for Disease Control (BCCDC), MSP billings, hospitalization data from the Discharge Abstract Database, the Sunquest Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory, Providence Health Care laboratory and the BC-CFE Drug Treatment Program (DTP) Database.

The objectives of these reports are to:

1. Provide timely HA-specific information on key HIV indicators which will guide and inform HIV leaders and innovators in the development of future HIV interventions and programs which will ultimately lead to decreasing the burden of HIV in BC. The indicators will reflect ongoing or past successful public health interventions and highlight areas in the HIV care spectrum which require further attention and support.
2. Highlight limitations in our current data due to incomplete or time lagged data and to develop future strategies to improve complete and timely data capture.

These reports are produced for the benefit of individual HAs. As such, we are enthusiastic about your involvement and cooperation regarding the development of these monitoring reports. Please forward your comments and queries to Irene Day, Director of Operations at the BC-CFE at iday@cfenet.ubc.ca.

** Please note that for Q2 and Q3 2013 reports, a coding revision resulted in data display errors in Indicator 5, the Cascade of Care (in Figures 5.1–5.7 on pp. 22–27 in this report), which has been updated; and, only for Q3 2013 reports, Appendix Table for Indicator 5 (on p. 37 in this report). All other figures and reports remain accurate. Please discard any previous reports and use this updated version. If you have any questions, please contact Irene Day at iday@cfenet.ubc.ca.*

List of Indicators

Indicator 1. Testing Episodes

Indicator 2. HIV Testing Rate

Indicator 3. New HIV Diagnoses

Indicator 4. Stage of HIV Infection at Diagnosis

Indicator 5. HIV Cascade of Care

Indicator 6. Programmatic Compliance Score (PCS)

Indicator 7. New Antiretroviral Starts

Indicator 8. CD4 Cell Count at ART Initiation

Indicator 9. Active and Inactive Drug Treatment Program Participants

Indicator 10. Antiretroviral Adherence Level

Indicator 11. Resistance Testing Results by Resistance Category

Indicator 12. AIDS-Defining Illness

Indicator 13. HIV-Related Mortality

Table of Contents

Acknowledgements and Contributions

BC Provincial STOP Program:

A Note on Monitoring and Interpreting HIV Indicators

Indicator 1 HIV Testing Episodes

- Figure 1.1 HIV Test Episodes for BC, 2009 Q1–2013 Q3
- Figure 1.2 HIV Test Episodes for BC by Gender and Prenatal Status,
2009 Q1–2013 Q3
- Figure 1.3 HIV Test Episodes for BC by Age Category, 2009 Q1–2013 Q3
- Figure 1.4 Point-of-Care HIV Tests for BC, 2010 Q4–2013 Q3
- Figure 1.5 HIV Test Episodes by Health Authority for BC, 2009 Q1–2013 Q3

Indicator 2 HIV Testing Rates

- Figure 2.1 Rate of HIV Testing for BC and Health Authorities, 2009–2012
- Figure 2.2 Rate of HIV Testing for BC by Gender, 2009–2012
- Figure 2.3 Rate of HIV Testing for BC by Age Category, 2009–2012

Indicator 3 New HIV Diagnoses

- Figure 3.1 New HIV Diagnoses for BC, 2009 Q1–2013 Q3
- Figure 3.2 New HIV Diagnoses by Gender for BC, 2009 Q1–2013 Q3
- Figure 3.3 New HIV Diagnoses by Age Category for BC, 2009 Q1–2013 Q3
- Figure 3.4 New HIV Diagnoses by Exposure Category for BC, 2009 Q1–2012 Q4
- Figure 3.5 New HIV Diagnoses by Health Authority for BC, 2009 Q1–2012 Q4

Indicator 4 Stage of HIV Infection at Diagnosis

- Table 1 Staging Classifications of Infection at Time of HIV Diagnosis Based on
CDC HIV Surveillance Case Definitions
- Figure 4.1 Stage of HIV Infection at Diagnosis for BC, 2010–2012
- Figure 4.2 Stage of HIV Infection at Diagnosis by Gender for BC, 2010–2012
- Figure 4.3 Stage of HIV Infection at Diagnosis by Age Category for BC, 2010–2012
- Figure 4.4 Stage of HIV Infection at Diagnosis by Exposure Category for BC,
2010–2012

Indicator 5 HIV Cascade of Care

- Figure 5.1 Estimated Cascade of Care for BC, Year Ending 2013 Q3
- Figure 5.2 Estimated Cascade of Care for BC by Gender, Year Ending 2013 Q3
- Figure 5.3 Estimated Cascade of Care for BC by Age Category,
Year Ending 2013 Q3
- Figure 5.4 Estimated Cascade of Care for BC by MSM Status, Year Ending 2013 Q3

Figure 5.5	Estimated Cascade of Care for BC by Age Category and MSM Status, Year Ending 2013 Q3
Figure 5.6	Estimated Cascade of Care for BC by History of IDU, Year Ending 2013 Q3
Figure 5.7	Estimated Cascade of Care for BC by Health Authority, Year Ending 2013 Q3
Indicator 6	Programmatic Compliance Score (PCS)
Table 2	Probability of Mortality Based on the Programmatic Compliance Score
Figure 6.1	PCS Components for BC, 2011 Q4–2013 Q3 <ul style="list-style-type: none"> First-Year CD4 Measurement First-Year VL measurement Baseline Resistance Testing Recommended Highly Active Antiretroviral Therapy (HAART) Baseline CD4 ≥ 200 cells/μL Suppression at 9 Months
Figure 6.2	Historical Trends for PCS Score for BC, 2011 Q4–2013 Q3
Indicator 7	New Antiretroviral Therapy Starts in BC
Figure 7	BC-CfE Drug Treatment Program Enrollment: New Antiretroviral Participants for BC, 2011 Q4–2013 Q3
Indicator 8	CD4 Cell Count at ART Initiation
Figure 8	CD4 Cell Count at ART Initiation for ART-Naïve DTP Participants for BC, 2011 Q4–2013 Q3
Indicator 9	Active and Inactive Drug Treatment Program (DTP) Participants
Table 3	Distribution of People on ART in BC by Health Authority, 2013 Q3
Figure 9	Active and Inactive DTP Participants for BC, 2011 Q4–2013 Q3
Indicator 10	Antiretroviral Adherence
Figure 10	Distribution of Individuals by Adherence Level in 1st Year of Therapy, Based on Pharmacy Refill Compliance for BC, 2011 Q4–2013 Q3
Indicator 11	Resistance Testing and Results
Figure 11	Cumulative Resistance Testing Results by Resistance Category for BC, 2011 Q4–2013 Q3
Indicator 12	AIDS-Defining Illness
Figure 12	AIDS Case Rate and Reports for BC, 2005–2012
Indicator 13	HIV-Related Mortality
Figure 13	HIV-Related Deaths by Year for BC, 2004–2011

Acknowledgements and Contributions



BRITISH COLUMBIA
CENTRE *for* EXCELLENCE
in HIV/AIDS

British Columbia Centre for Excellence in HIV/AIDS (BC-CFE): The BC-CFE is responsible for the conception, preparation and ongoing review of this quarterly report. The BC-CFE provides the data and outputs for Indicators 5 (Hiv Cascade of Care), 6 (Programmatic Compliance Score), 7 (New Antiretroviral Starts), 8 (CD4 Cell Count at ART Initiation), 9 (Active and Inactive Drug Treatment Program Participants), 10 (Antiretroviral Adherence Level), 11 (Resistance Testing Results by Resistance Category), 12 (AIDS-Defining Illness), and 13 (HIV-Related Mortality). The BC-CFE database provides PVL and CD4 cell count testing data, as well as ART use. All PVL measurements in BC are performed at the St Paul's Hospital virology laboratory, thus PVL data capture is 100%. An estimated 80% of all CD4 count measurements performed in the province are captured in the BC-CFE data holdings. The STOP HIV/AIDS Technical Monitoring Committee-BC-CFE is responsible for oversight of the monitoring report. Lillian Lourenco writes and compiles the monitoring report. Guillaume Colley, Dr. Viviane Lima and Nada Gataric perform analysis of Indicators 5–13. James Nakagawa is responsible for publishing and editing. This report was conceived and guided by Dr. Julio Montaner.



BC Centre for Disease Control
An agency of the Provincial Health Services Authority

British Columbia Centre for Disease Control (BCCDC): The BCCDC provides the data and outputs for Indicator 1 (Hiv Testing Episodes), Indicator 2 (Hiv Testing Rate), Indicator 3 (New Hiv Diagnoses), Indicator 4 (Stage of Hiv at Diagnosis) and Indicator 12 (AIDS-Defining Illness). The BCCDC is the single provincial agency that centralizes all HIV surveillance through the Public Health Microbiology and Reference Laboratory, which does more than 90% of all HIV screening tests in BC and all confirmatory testing. Theodora Consolacion and Dr. Mark Gilbert are responsible for outputs for Indicators 1–4.

Other Data Sources:

The above databases were supplemented with:

- (I) The BC Vital Statistics database which was used to calculate Indicator 5. The Hiv Cascade of Care and Indicator 13. HIV-Related Mortality.
- (II) Linkage and preparation of the de-identified individual-level database used for calculating Indicator 5. The Hiv Cascade of Care was facilitated by the British Columbia Ministry of Health.
- (III) The Statistics Canada database: BC and HIV-positive population counts were acquired through the statistics Canada website to calculate HIV-specific mortality rates for Indicator 13. HIV-Related Mortality.

Membership of the STOP HIV/AIDS Technical Monitoring Committee–BC-CfE

Dr. Rolando Barrios, *Chair*, BC-CfE

Kate Heath, BC-CfE

Bohdan Nosyk, BC-CfE

Viviane Dias Lima, BC-CfE

Irene Day, BC-CfE

Dr. Mark Gilbert, BCCDC

Dr. Mel Kradjen, BCCDC

Stephanie Konrad, FHA

Joanne Nelson, FNHA

Jennifer May-Hadford, IHA

James Haggerstone, NHA

Dr. Neora Pick, PHSA

Dr. Reka Gustafson, VCHA

Melanie Rusch, VIHA

The Seek and Treat for Optimal Prevention (STOP) HIV/AIDS BC Provincial Program: A Note on Monitoring and Interpreting HIV Indicators

The Seek and Treat for Optimal Prevention (STOP) of HIV/AIDS programme is a provincial initiative to improve HIV diagnosis and care delivery in BC through increased HIV-specific funding to all HSDA's across BC. The STOP provincial programme is an expansion of a four-year STOP pilot project which was implemented in two Health Service Delivery Areas in March 2010; the Vancouver HSDA which bears the largest burden of the HIV epidemic in the province and the Northern Interior HSDA which bears a high burden of HIV-related mortality. The STOP pilot project demonstrated the urgent need for improved efforts in early diagnosis of HIV and timely initiation of highly active antiretroviral therapy (HAART) initiation.

The expansion to a province-wide programme was announced on November 30th 2012 by the BC Ministry of Health with roll out of funding beginning on April 1st, 2013. This funding is intended to be used in the implementation and evaluation of HIV-related diagnosis and care initiatives within individual HA's. Goals of the project include: 1. A reduction in the number of new HIV infections in BC; 2. Improvements in the quality, effectiveness, and reach of HIV prevention services; 3. An increase in early diagnosis of HIV; 4. A reduction in AIDS cases and HIV-related mortality.

The goals of HA-led STOP-funded initiatives are to work toward achieving these goals. To these ends some outcome measures or indicators of progress have been drafted that should be considered in the design and implementation phases of these initiatives.

HIV Testing Episodes and Rates

In this section, the number of HIV test episodes and point of care (POC) HIV tests conducted each quarter in BC is shown. In general terms the goal is to increase the number of tests performed and to maximize testing efficiency. Test episodes are allocated by region according to where the test is performed.

NB: HIV screening tests conducted by the VIHA Laboratory are not included.

Indicator 1. HIV Testing Episodes

Figure 1.1 HIV Test Episodes in BC, 2009 Q1–2013 Q3





Figure 1.2 HIV Test Episodes by Gender and Prenatal Status in BC, 2009 Q1–2013 Q3¹




1 NB: Testing does not include point of care tests.

Figure 1.3 HIV Test Episodes by Age Category for BC, 2009 Q1–2013 Q3 ^{1,2}



Figure 1.4 Point-of-Care HIV Tests in BC, 2010 Q4–2013 Q3




² Data Source: The BC Public Health Microbiology and Reference Laboratory (BCPHMRL) courtesy of the BC Centre for Disease Control (BCCDC).

Limitations:

- 1 Repeat tests in individuals who test using various identifiers may not be identified and these individuals may be counted more than once.
- 2 POC testing data is available from the fourth quarter of 2010 and onwards.

Figure 1.5 HIV Test Episodes by Health Authority in BC, 2009 Q1–2013 Q3



Indicator 2. HIV Testing Rates

Figure 2.1 Rate of HIV Testing in BC and Health Authorities, 2009–2012 ¹




Figure 2.2 Rate of HIV Testing by Gender in BC, 2009–2012 ¹





Figure 2.3 Rate of HIV Testing by Age Category in BC, 2009–2012 ¹



New HIV Diagnoses

Trends in HIV diagnoses by gender and exposure category are described. Interpreting HIV diagnoses must be done with consideration that trends are influenced by both changes in testing rate as well as changes in transmission rates. It is important to note that new HIV diagnoses cases and rates are not synonymous with HIV incidence as a person may have become infected with HIV long before they tested positive for HIV. However, as there is no reliable method for measuring HIV incidence we follow trends in HIV diagnoses.

Indicator 3. New HIV Diagnoses

Figure 3.1 New HIV Diagnoses for BC, 2009 Q1–2013 Q3 ³




Figure 3.2 New HIV Diagnoses for BC by Gender, 2009 Q1–2013 Q3




Figure 3.3 New HIV Diagnoses for BC by Age Category, 2009 Q1–2013 Q3






Figure 3.4 New HIV Diagnoses for BC by Exposure Category, 2009 Q1–2013 Q3⁴



⁴ BCCDC: Data lags by 6 months.

MSM=men who have sex with men; IDU= injection drug user; HET=heterosexual. NIR=No identified risk/exposure.

Figure 3.5 New HIV Diagnoses for BC by Health Authority, 2009 Q1–2013 Q3



Fraser Health	By Client Residence	27	27	21	12	17	19	19	17	12	18	15	9	11	10	10	14	9	16	22
	By Provider Address	22	17	17	7	14	15	13	15	10	20	9	8	10	5	7	8	9	9	14
Interior Health	By Client Residence	6	4	4	4	4	1	3	3	1	4	3	3	5	5	0	2	1	4	5
	By Provider Address	6	4	4	4	4	1	2	3	1	3	2	3	5	5	0	2	1	5	5
Northern Health	By Client Residence	7	8	5	7	4	3	7	2	4	10	5	5	5	4	3	0	3	8	3
	By Provider Address	7	8	5	8	4	3	6	2	4	9	5	6	5	4	3	0	3	8	3
Vancouver Coastal Health	By Client Residence	51	36	41	36	41	45	41	40	32	48	57	41	43	33	34	32	32	45	50
	By Provider Address	56	46	45	40	45	50	49	43	34	47	65	42	47	38	38	38	32	51	58
Vancouver Island Health	By Client Residence	9	9	7	14	6	13	8	6	5	2	7	7	6	3	5	12	4	10	12
	By Provider Address	11	9	7	14	6	12	8	5	5	3	6	6	3	3	5	12	5	10	12

"By Provider Address" is graphed as dashed line in same colour.

Stage of HIV infection at diagnosis

Classification of stage of HIV infection, in the absence of information regarding recent testing history, is reliant on clinical information available at the time of diagnosis, including first CD4+ cell count, laboratory results suggestive of acute HIV infection, and clinical presentation with an AIDS-defining illness (Table 1). The benefits of Treatment as Prevention (TasP) are maximized when antiretroviral therapy (ART) is initiated at high CD4 cell counts. Accordingly, it is preferable that individuals newly diagnosed with HIV be in the early stages of HIV infection (stage 0 or 1) to allow for early ART initiation.

N.B. Interpretation of stage of HIV infection at diagnosis should proceed with caution. Early increases in diagnosis at late stage (i.e., low CD4 counts) may represent a “catching up” of previously missed long term infected individuals rather than a trend toward diagnosis at later stage of infection.

Indicator 4. Stage of HIV Infection at Diagnosis

Table 1 Staging Classifications of Infection at Time of HIV Diagnosis Based on CDC HIV Surveillance Case Definitions

Stage	Criteria		
0	Laboratory criteria met for acute HIV infection, or previous negative or indeterminate HIV test within 180 days of first confirmed positive HIV test.		
1	Stage 0 not met	CD4 ≥500	and No AIDS case report
2a		CD4 350–499	
2b		CD4 200–349	
3		(CD4 <200 or AIDS case report)	
Unknown		No available CD4	and No AIDS case report

Figure 4.1 Stage of HIV Infection at Diagnosis for BC, 2010–2012 ⁵





Figure 4.2 Stage of HIV Infection at Diagnosis by Gender for BC, 2010–2012 ⁵



	BC			Female			Male		
	2010	2011	2012	2010	2011	2012	2010	2011	2012
Stage 0	48	63	53	8	3	6	40	60	47
Stage 1	67	52	52	12	7	6	55	45	46
Stage 2a	35	39	28	5	7	5	30	32	23
Stage 2b	36	39	31	12	8	6	24	31	25
Stage 3	76	69	57	22	13	5	54	56	52
Unknown	38	26	17	3	5	1	35	21	16
Total (n=)	300	288	238	62	43	29	238	245	209

5 Data Source: BCCDC

Figure 4.3 Stage of HIV Infection at Diagnosis by Age Category for BC, 2010–2012 ⁵





Figure 4.4 Stage of HIV Infection at Diagnosis by Exposure Category for BC, 2010–2012 ^{5,6}



	< 30 years			30–39 years			40–49 years			≥ 50 years			MSM			HET			IDU			Other			NIR/Unknown		
	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012
Stage 0	16	14	18	15	25	18	13	19	10	4	5	7	34	51	42	7	7	7	6	5	4	0	0	0	1	0	0
Stage 1	20	12	11	20	20	14	20	9	16	7	11	11	34	31	33	20	13	12	13	8	6	0	0	1	0	0	0
Stage 2a	8	11	4	14	12	4	9	11	11	4	5	9	21	25	17	8	7	5	6	5	6	0	2	0	0	0	0
Stage 2b	6	6	7	10	8	8	11	12	10	9	13	6	16	20	17	13	15	8	5	3	6	2	1	0	0	0	0
Stage 3	2	5	7	13	17	10	31	26	15	30	21	25	25	28	29	28	31	20	18	7	6	3	3	0	2	0	2
Unknown	11	6	7	15	10	3	11	5	2	1	5	5	23	15	11	7	2	1	4	6	1	1	1	1	3	2	3
Total (n=)	63	54	54	87	92	57	95	82	64	55	60	63	153	170	149	83	75	53	52	34	29	6	7	2	6	2	5

6 MSM=men who have sex with men; IDU= injection drug user; HET=heterosexual. NIR=No identified risk/exposure.

Indicator 5. HIV Cascade of Care

The success of seek, test, treat and retain (STTR) strategies like STOP is reliant on early diagnosis of HIV, linking newly diagnosed HIV-positive persons with ongoing care, retaining persons in HIV-care; initiating ART based on best evidenced practices and maintaining optimal ART adherence to ensure a suppressed viral load. These stages of HIV-care can be summarized as: 1. HIV diagnosis, 2. Linkage to HIV care, 3. Retention in HIV care, 4. On ART and 5. Achieving a suppressed VL; collectively, they are referred to as the cascade of care. Leakage between any of these stages of HIV-care means a reduction in the potential of ART as a benefit to the HIV-positive individual and as an HIV transmission prevention method on a population level. Thus, when interpreting trends in the cascade of care, we strive to see increases along each step of the cascade of care (ie. reduced attrition) with the ultimate goal being 100% within each stage of the cascade. Monitoring the Cascade of Care provides a picture as to where deficiencies lie in the delivery and uptake of HIV-care. In this section we present the cascade of care for the year 2012 in BC overall and stratified by sex and age for each Health Authority.

Figure 5.1 Estimated Cascade of Care for BC, Year Ending 2013 Q3 ⁷





Figure 5.2 Estimated Cascade of Care for British Columbia by Gender, Year Ending 2013 Q3 ⁸




7,8 Data Sources:

- 1 British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- 2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).


NB: Transgender has been assigned to their biological sex.

Figure 5.3 Estimated Cascade of Care for British Columbia by Age Category, Year Ending 2013 Q3 ⁹



⁹ Data Sources:
1 British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).


Figure 5.4 Estimated Cascade of Care for British Columbia by MSM Status, Year Ending 2013 Q3 ¹⁰



¹⁰ Data Sources:

- 1 British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- 2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).


Figure 5.5 Estimated Cascade of Care for British Columbia by Age Category and MSM Status, Year Ending 2013 Q3 ¹¹



¹¹ Data Sources:

- 1 British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- 2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).


Figure 5.6 Estimated Cascade of Care for British Columbia by History of IDU, Year Ending 2013 Q3 ¹²



¹² Data Sources:

- ¹ British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- ² Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Figure 5.7 Estimated Cascade of Care for British Columbia by Health Authority, Year Ending 2013 Q3 ¹³



¹³ Data Sources:

- 1 British Columbia Centre for Excellence Drug Treatment Program (dtp) Database (ARV use, VL and CD4 count).
- 2 Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider. If the most recent HA of residence is not updated then the designated HA may be incorrect.

Indicator 6. The Programmatic Compliance Score (PCS)

The Programmatic Compliance Score (PCS) is a summary measure of risk of future death, immunologic failure and virologic failure from all causes for people who are starting ART for the first time. It is composed of patient- and physician-driven effects. PCS scores range from 0–6 with higher scores indicative of poorer health outcomes and greater risk of death. Table 2 provides mortality, immunologic failure and virologic failure probabilities for given PCS scores. We interpret an individual with a $PCS \geq 4$ as being 22 times more likely to die, almost 10 times more likely to have immunologic failure and nearly 4 times as likely to demonstrate virologic failure compared to those individuals with a PCS score of 0. A detailed description of how the PCS score is calculated and its validation can be found in the technical report. In short, PCS scores are calculated by summing the results (yes=1, no=0) of six un-weighted non-performance indicators based on IAS–USA treatment guidelines:

1. having <3 CD4 cell count tests in the first year after starting antiretroviral therapy (ART);
2. having <3 plasma viral load (VL) tests in the first year after starting ART;
3. not having drug resistance testing done prior to starting ART;
4. starting on a non-recommended ART regimen;
5. starting therapy with $CD4 < 200$ cells/ μ L; and
6. not achieving viral suppression within 9 months since ART initiation.

In this section we provide PCS scores and their components over time for the province of BC. A decline to 0%, (i.e., **all individuals having a score of 0**) is the eventual goal.

Table 2. The Probability of Mortality, Immunologic Failure and Virologic Failure based on the Programmatic Compliance Score

Programmatic Compliance Score	Mortality Risk Ratio (95% Confidence Interval)	Immunologic Failure Risk Ratio (95% CI)	Virologic Failure Risk Ratio (95% CI)
0 (Best score)	1 (–)	1 (–)	1 (–)
1	3.81 (1.73–8.42)	1.39 (1.04–1.85)	1.32 (1.05–1.67)
2	7.97 (3.70–17.18)	2.17 (1.54–3.04)	1.86 (1.46–2.38)
3	11.51 (5.28–25.08)	2.93 (1.89–4.54)	2.98 (2.16–4.11)
4 or more (Worst score)	22.37 (10.46–47.84)	9.71 (5.72–16.47)	3.80 (2.52–5.73)

Reference: Lima VD, Le A, Nosyk B, Barrios R, Yip B, et al. (2012) Development and Validation of a Composite Programmatic Assessment Tool for HIV Therapy. *PLoS ONE* 7(11): e47859. doi:10.1371/journal.pone.0047859

Figure 6.1 PCS Components for BC, 2011 Q4–2013 Q3 ¹⁴





Figure 6.2 Historical Trends for PCS Score for BC, 2011 Q4–2013 Q3 ¹⁵



NB: A score of 0 is the best score and a score of 4 or more is the worst score.

¹⁴ Data Source: British Columbia Centre for Excellence Drug Treatment Program (DTP) Database.
Limitations: CD4 cell count capture is approximately 80%.

¹⁵ Data Source: British Columbia Centre for Excellence Drug Treatment Program (DTP) Database.
Each quarter's data is calculated as the sum of the 4 quarters leading up to it. e.g. 2012 Q1 is calculated from 2011 Q2 – 2012 Q1.


Antiretroviral Uptake

In this section we present trends in ART uptake, the number and proportion of new HIV treatment initiations and the number of active and inactive DTP participants. Trends in ART uptake should be interpreted under the consideration of changing BC HIV treatment guidelines. BC HIV treatment guidelines are updated regularly by the BC-CfE Therapeutic Guidelines Committee and reflect those of the International AIDS Society. Most recent changes were made in 2012 and HIV treatment is now recommended for all HIV-positive adults regardless of CD4 cell count; as evidence demonstrates that early initiation of HIV treatment maximizes both the individual's health outcomes as well as the potential of ART as a form of HIV transmission prevention at a population level. As such, trends in the number and proportion of persons on ART and new ART starts (in both naïve and experienced persons) are expected to increase over time at higher CD4 cell counts.

Indicator 7. New Antiretroviral Therapy Starts in BC

Figure 7


BC-CfE Drug Treatment Program Enrollment: New Antiretroviral Participants for BC, 2011 Q4–2013 Q3¹⁶



Indicator 8. CD4 Cell Count at ART Initiation

Figure 8

CD4 Cell Count at ART Initiation of ART-Naïve DTP Participants for BC, 2011 Q4–2013 Q3¹⁷



¹⁶ Data Source: Drug Treatment Program Database

¹⁷ Data Source: Drug Treatment Program Database


Limitations: CD4 cell count data is approximately 80% complete.

Indicator 9. Active and Inactive DTP Participants

Table 3. Distribution of People on ART in BC by Health Authority, 2013 Q3 ¹⁸

		Interior	Fraser	Vancouver Coastal	Vancouver Island	Northern	Total BC
Age	< 30	9	64	141	29	15	258
	30–39	52	259	508	96	43	958
	40–49	113	516	1256	222	61	2168
	≥ 50	239	702	1838	410	81	3271
Gender	Male	321	1191	3293	615	117	5538
	Female	92	350	450	142	83	1117
Exposure	MSM	112	451	1587	180	20	2350
	IDU	144	456	1094	272	122	2088
Total		413	1541	3743	757	200	6655

Figure 9 Active and Inactive DTP Participants for BC, 2011 Q4–2013 Q3 ¹⁹



¹⁸ Data Source: Drug Treatment Program Database

Limitation: DTP participants are designated to an HA based on most current residence provided by the participant.

Definitions:

'On antiretroviral therapy' defined as being on treatment in the current quarter

'Unknown/not stated' defined as being on treatment in the current quarter, and city of residence unknown

¹⁹ Active DTP participants: are those who are prescribed one or more drugs in the last six months.


Inactive DTP Participants: Persons no longer prescribed drugs through the HIV/AIDS Drug Treatment Program in the last quarter.

Antiretroviral Adherence Level

In this section we present trends in prescription refill adherence levels for individuals in their first year of treatment. Given that the benefits of ART are compromised in the presence of imperfect ART adherence, we expect to see the proportion of persons on ART achieving **near perfect adherence** (ie. $\geq 95\%$) to increase with time. Furthermore, it is important that trends in the proportion of ART users achieving prescription refill adherence of $\geq 95\%$ keep pace with new ART starts and increase among those continuing on ART.

Indicator 10. Antiretroviral Adherence


Figure 10 Distribution of Individuals by Adherence Level in 1st Year of Therapy, Based on Pharmacy Refill Compliance for BC, 2011 Q4–2013 Q3²⁰



²⁰ Data Source: Drug Treatment Program Database
Limitation: Prescription refill adherence is used as a proxy for patient adherence.

Indicator 11. Resistance Testing and Results

In this section, we present trends in cumulative resistance testing by resistance category: **Suppressed** (where a DTP participant's viral load is too low to be genotyped); **Wild Type** (where no HIV treatment resistances were discovered), **Never Genotyped**, and Resistances to **one**, **two** or **three** HIV treatment classes. Resistance testing prior to ART initiation is recommended in the BC HIV treatment primary care guidelines. Thus, it is expected that trends over time should find all persons enrolled in the DTP to have been genotyped. Trends over time should also show an increase in the proportion of DTP participants achieving a suppressed status and an increase in resistance testing should not lead to an increase in the number of ART resistances occurring.




²¹ Data Source: Drug Treatment Program Database

Indicator 12. AIDS-Defining Illness

Improvements in ART and the expansion of ART province-wide has led to very low numbers of recorded AIDS cases across BC. However, interpreting trends in AIDS cases is challenging as AIDS reporting is passive in BC and it is likely that they are under reported across all Health Authorities. In addition to under reporting, methods of reporting AIDS cases are inconsistent across HA's and do not truly reflect the current reality of new AIDS diagnoses. Efforts will need to be made to improve under and inconsistent reporting of AIDS cases across all HA's. The table below shows AIDS cases using three definitions. First, AIDS cases were defined as the number of physician-reported AIDS defining illness (ADI) in a given year. AIDS case reporting is a passive process and physicians can voluntarily report AIDS cases to the BCCDC or DTP. As such, we have plotted both **BCCDC reports** and **DTP reported** AIDS cases. We also show the proportion of persons **initiating ART with a CD4<200 cells/μL**.

Figure 12 AIDS Case Rate and Reports for BC, 2005–2012 ²²




²² Data Source: DTP AIDS cases are obtained from the Drug Treatment Program Database; BCCDC AIDS cases are obtained from the BC-CDC; CD4<200 at ART initiation data came from the DTP database.

Limitation: AIDS case reporting was investigated using 3 definitions: First, using AIDS cases reported in AIDS case report forms from the DTP; Second, using AIDS cases reported via the BCCDC and third, using a CD4 cell count of <200 cells/μL at time of ART initiation using DTP data. AIDS case reporting is passive in BC, thus; AIDS case reporting is not well captured. The DTP sends out AIDS reporting forms to physicians annually. The BCCDC uses DTP AIDS case reports as well as physician AIDS case reports made directly to the BCCDC. Interpreting AIDS case reports should be done with these limitations in mind. AIDS data is updated annually as very few AIDS cases reports are reported in general and trends would be difficult to notice if reported quarterly.

Indicator 13. HIV-Related Mortality

Evidence indicates that individuals who initiate treatment with recommended ART in a timely fashion may live near normal lifespans. Excess mortality among HIV positive persons is, therefore, an important measure of HIV care with a goal of minimizing HIV-related mortality in British Columbia.

Figure 13 HIV-Related Deaths by Year for BC, 2004–2011 ²³



²³ Data Source: BC Vital Statistics
Limitation: Mortality data is updated annually. The most recent available data was used.

Appendices

Indicator 1: Test Episodes (thousands)		2009				2010				2011				2012				2013			
		Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	
B.C. Overall		47.2	43.6	44.0	41.5	46.3	44.0	44.6	45.4	48.6	45.4	52.9	50.8	56.7	55.2	59.2	59.8	67.2	69.7	67.4	
Gender	Female	28.4	26.4	26.6	25.5	28.0	26.3	27.0	26.9	28.7	26.2	28.0	29.0	32.4	31.6	33.8	33.9	38.1	39.2	38.3	
	Male	18.1	16.5	16.7	15.4	17.6	17.1	16.9	17.0	18.2	16.6	18.2	18.8	21.6	21.1	22.8	23.4	26.2	27.8	26.4	
	Other	0.7	0.7	0.7	0.6	0.7	0.7	0.6	0.6	0.6	0.6	0.6	0.6	0.4	0.4	0.4	0.3	0.3	0.3	0.3	
Female (Prenatal)		12.7	11.2	11.4	11.3	12.1	11.0	11.8	12.0	12.9	11.3	11.8	11.9	13.0	11.9	12.0	11.9	12.5	11.7	12.0	
Female (Non-prenatal)		15.8	15.2	15.2	14.1	15.9	15.2	15.3	14.9	15.8	14.9	16.2	17.1	19.4	19.7	21.8	22.0	25.6	27.5	26.3	
Age	< 30	19.2	18.0	19.0	17.4	18.5	17.9	18.5	18.5	18.5	17.5	19.2	19.9	20.5	20.1	21.7	21.2	20.6	21.2	21.6	
	30–39	15.3	14.0	13.8	13.2	15.4	14.2	14.4	14.1	15.9	13.9	14.8	14.8	17.3	16.3	16.8	16.6	19.2	19.1	18.8	
	40–49	7.0	6.4	6.2	5.7	6.7	6.4	6.1	6.2	6.8	6.3	6.5	6.6	7.8	7.5	7.7	8.2	9.7	10.1	9.4	
	≥ 50	5.7	5.2	5.1	5.2	5.7	5.6	5.5	5.7	6.2	5.7	6.3	7.0	8.8	9.1	10.8	11.7	15.2	16.9	15.2	
POC HIV Tests									1.0	1.1	2.1	6.1	2.5	2.3	2.1	2.3	2.2	2.6	2.4	2.4	
Fraser Health		13.0	12.4	12.3	11.6	13.0	12.3	12.5	12.4	13.1	12.2	13.0	13.0	14.4	14.2	15.1	15.6	17.9	18.4	18.2	
Interior Health		6.0	5.6	5.5	5.6	6.0	5.6	5.5	5.9	6.0	5.5	5.6	5.9	5.9	5.7	5.9	5.9	6.1	6.4	6.3	
Northern Health		3.1	2.6	2.7	2.5	2.9	2.6	2.7	2.5	3.0	2.7	2.9	2.8	3.4	3.1	3.2	3.3	3.7	3.6	3.5	
Van. Coastal Health		18.9	17.9	18.3	17.1	18.8	18.3	18.6	19.3	20.7	20.1	26.0	24.0	27.2	26.9	29.5	29.4	33.5	35.1	33.4	
Van. Island Health		6.2	5.1	5.2	4.7	5.6	5.3	5.3	5.3	5.8	5.0	5.3	5.1	5.8	5.3	5.5	5.6	5.9	6.2	6.0	

Indicator 2: **Rate of HIV Testing per 100,000**

		2009	2010	2011	2012
British Columbia Overall		3 633.4	3 627.7	3 717.0	4 380.4
Fraser Health		3 385.8	3 381.3	3 439.2	3 895.1
Interior Health		2 890.4	2 926.0	2 929.2	2 967.9
Northern Health		3 524.4	3 442.8	3 592.8	4 100.0
Vancouver Coastal Health		5 114.6	5 088.5	5 342.2	7 111.4
Vancouver Island Health		2 720.4	2 702.2	2 659.9	2 769.6
Gender	Female	4 401.4	4 370.4	4 467.0	5 194.4
	Male	2 732.0	2 762.3	2 851.1	3 489.1
Age	< 30	4 292.7	4 222.3	4 303.5	4 766.2
	30–39	8 881.6	9 096.9	9 319.9	10 234.2
	40–49	3 351.6	3 357.9	3 479.7	4 207.1
	≥ 50	1 241.3	1 279.5	1 390.0	2 167.0

		2009				2010				2011				2012				2013			
Indicator 3: New HIV Diagnoses		Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	
B.C. Overall		102	84	78	73	73	81	78	68	54	82	87	65	70	55	53	60	50	83	92	
Gender	Female	21	16	17	17	17	20	14	11	6	19	11	7	9	10	5	5	6	12	14	
	Male	81	68	61	56	56	61	64	57	48	63	76	58	61	45	48	55	44	71	78	
Age	< 30	18	16	16	24	17	20	22	9	5	18	17	18	18	14	8	18	9	18	24	
	30–39	26	27	17	20	25	18	23	20	18	30	30	13	16	17	11	10	17	25	20	
	40–49	35	25	30	21	23	29	19	24	18	22	22	19	20	11	19	19	12	14	23	
	≥ 50	23	16	15	8	8	14	14	15	13	12	18	15	16	13	15	13	12	26	25	
Exposure	MSM	45	43	33	32	39	36	45	33	33	43	56	38	42	34	38	36	28	–	–	
	IDU	22	15	13	14	14	16	12	10	8	11	12	3	14	7	2	6	3	–	–	
	HET	28	21	21	22	18	25	18	22	12	23	19	21	14	12	11	15	9	–	–	
	Other	4	2	3	3	2	2	1	1	1	3	0	3	0	1	0	1	1	–	–	
	NIR	3	3	8	2	0	2	2	2	0	2	0	0	0	1	2	2	9	–	–	
Fraser Health	By Client Residence	27	27	21	12	17	19	19	17	12	18	15	9	11	10	10	14	9	16	22	
	By Provider Address	22	17	17	7	14	15	13	15	10	20	9	8	10	5	7	8	9	9	14	
Interior Health	By Client Residence	6	4	4	4	4	1	3	3	1	4	3	3	5	5	0	2	1	4	5	
	By Provider Address	6	4	4	4	4	1	2	3	1	3	2	3	5	5	0	2	1	5	5	
Northern Health	By Client Residence	7	8	5	7	4	3	7	2	4	10	5	5	5	4	3	0	3	8	3	
	By Provider Address	7	8	5	8	4	3	6	2	4	9	5	6	5	4	3	0	3	8	3	
Vancouver Coastal Health	By Client Residence	51	36	41	36	41	45	41	40	32	48	57	41	43	33	34	32	32	45	50	
	By Provider Address	56	46	45	40	45	50	49	43	34	47	65	42	47	38	38	38	32	51	58	
Vancouver Island Health	By Client Residence	9	9	7	14	6	13	8	6	5	2	7	7	6	3	5	12	4	10	12	
	By Provider Address	11	9	7	14	6	12	8	5	5	3	6	6	3	3	5	12	5	10	12	

Indicator 4: Stage of HIV Infection at Baseline

	BC			Female			Male			< 30 years			30–39 years			40–49 years		
	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012
Stage 0	48	63	53	8	3	6	40	60	47	16	14	18	15	25	18	13	19	10
Stage 1	67	52	52	12	7	6	55	45	46	20	12	11	20	20	14	20	9	16
Stage 2a	35	39	28	5	7	5	30	32	23	8	11	4	14	12	4	9	11	11
Stage 2b	36	39	31	12	8	6	24	31	25	6	6	7	10	8	8	11	12	10
Stage 3	76	69	57	22	13	5	54	56	52	2	5	7	13	17	10	31	26	15
Unknown	38	26	17	3	5	1	35	21	16	11	6	7	15	10	3	11	5	2
Total	300	288	238	62	43	29	238	245	209	63	54	54	87	92	57	95	82	64
	≥ 50 years			MSM			Heterosexual			IDU			Other Exposure			NIR/Unknown		
	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012	2010	2011	2012
Stage 0	4	5	7	34	51	42	7	7	7	6	5	4	0	0	0	1	0	0
Stage 1	7	11	11	34	31	33	20	13	12	13	8	6	0	0	1	0	0	0
Stage 2a	4	5	9	21	25	17	8	7	5	6	5	6	0	2	0	0	0	0
Stage 2b	9	13	6	16	20	17	13	15	8	5	3	6	2	1	0	0	0	0
Stage 3	30	21	25	25	28	29	28	31	20	18	7	6	3	3	0	2	0	2
Unknown	1	5	5	23	15	11	7	2	1	4	6	1	1	1	1	3	2	3
Total	55	60	63	153	170	149	83	75	53	52	34	29	6	7	2	6	2	5

Indicator 5: HIV Cascade of Care		DIAGNOSED		LINKED		RETAINED		ON ART		ADHERENT		SUPPRESSED	
B.C. Overall		9284		8338		7123		6541		6027		4899	
Age Category	< 30	476		324		265		226		189		151	
	30–39	1350		1189		1001		865		767		606	
	40–49	2962		2700		2279		2091		1894		1508	
	≥ 50	4488		4125		3579		3359		3177		2634	
Age Category and MSM Status	MSM	< 30		77		71		56		48		39	
		30–39		283		267		235		209		158	
		40–49		800		776		716		680		520	
		≥ 50		1604		1564		1510		1445		1202	
	Non-MSM	< 30		67		65		60		49		27	
		30–39		430		409		372		321		189	
		40–49		935		898		845		771		482	
		≥ 50		1271		1226		1164		1090		774	
	Unknown	< 30		333		187		149		129		85	
		30–39		637		514		394		335		259	
		40–49		1228		1026		719		640		506	
		≥ 50		1612		1335		904		824		658	
Gender	Male	7589		6869		5877		5441		5093		4184	
	Female	1695		1469		1246		1100		934		715	
Injection Drug Use	IDU	2526		2432		2323		2129		1870		1390	
	Non-IDU	3938		3817		3586		3394		3211		2728	
	Unknown	2820		2089		1214		1018		946		781	
MSM Status	MSM	2763		2679		2517		2382		2252		1919	
	Non-MSM	2704		2597		2440		2231		1966		1472	
	Unknown	3817		3062		2166		1928		1809		1508	
Health Authority	Fraser Health	1885		1788		1650		1528		1399		1159	
	Interior Health	535		514		440		407		380		287	
	Island Health	886		847		789		740		666		496	
	Northern Health	267		247		217		192		159		104	
	Vancouver Coastal Health	4487		4283		3924		3673		3422		2852	

Indicator 6: Programmatic Compliance Score (PCS)

	2011 Q4	2012 Q1	Q2	Q3	Q4	2013 Q1	Q2	Q3
< 3 CD4 Tests	10.2%	10.0%	9.2%	8.1%	8.6%	7.0%	7.5%	9.5%
< 3 Viral Load Tests	6.3%	7.8%	7.5%	7.0%	7.6%	5.0%	6.0%	6.7%
No Baseline Genotype	6.6%	6.1%	6.4%	6.0%	5.7%	5.5%	5.0%	5.0%
Baseline CD4 < 200 cells/μL	30.0%	29.0%	26.9%	25.3%	23.6%	22.2%	22.6%	23.7%
Non-Recommended ART	2.4%	2.8%	2.5%	2.3%	2.2%	1.5%	2.0%	5.8%
Non Viral suppression at 9 Mo.	26.1%	25.3%	23.3%	24.7%	26.4%	24.7%	25.6%	26.2%
PCS Score: 0	165	180	188	202	212	222	220	184
PCS Score: 1	94	104	105	115	118	113	103	98
PCS Score: 2	52	48	42	42	51	49	59	57
PCS Score: 3	15	19	18	19	20	15	15	16
PCS Score: 4 or more	7	8	7	6	5	2	2	4
Total (n=)	333	359	360	384	406	401	399	359

**Indicator 7: New DTP
ARV Participants**

	2011 Q4	2012 Q1	Q2	Q3	Q4	2013 Q1	Q2	Q3
First Starts	117	129	105	94	92	97	94	113
Experienced Starts	92	96	106	89	109	91	111	113

Indicator 8: CD4 Cell Count at ART Initiation for ARV-Naïve DTP Participants

CD4 ≥ 500	28	36	36	24	35	21	36	32
CD4 350–499	36	37	26	30	21	29	16	20
CD4 200–349	28	32	22	15	15	25	19	26
CD4 50–199	16	12	12	18	12	15	13	23
CD4 < 50	8	11	9	7	7	7	8	10
<i>CD4 Median (cells/μL)</i>	<i>380</i>	<i>368</i>	<i>390</i>	<i>370</i>	<i>430</i>	<i>360</i>	<i>381</i>	<i>300</i>
Total (n=)	116	128	105	94	90	97	92	111

Indicator 9: Active and Inactive DTP Participants

Active DTP Participants	6155	6258	6342	6403	6459	6497	6581	6653
Inactive DTP Participants	1217	1200	1198	1219	1223	1251	1235	1249

Indicator 10: Antiretroviral Adherence

≥ 95%	70	105	90	100	77	98	88	64
80% to < 95%	22	19	23	28	23	21	20	23
40% to < 80%	14	10	11	9	14	7	9	8
< 40%	2	2	0	3	0	2	1	1
Total (n=)	108	136	124	140	114	128	118	96

Indicator 11: Resistance Testing and Results

Suppressed	4054	4319	4337	4378	4240	4563	4584	4665
Wild Type	1067	1048	979	939	984	947	888	846
Never Genotyped	100	109	85	102	103	81	92	101
1-Class	203	221	197	186	193	173	185	168
2-Class	42	37	43	38	45	43	48	37
3-Class	5	7	3	10	7	9	9	10
Total (n=)	5471	5741	5644	5653	5572	5816	5806	5827

Indicator 12: AIDS-Defining Illness

	2005	2006	2007	2008	2009	2010	2011	2012
CD4 < 200 at Cases	217	204	219	194	153	133	112	90
ART initiation <i>Rate per 100,000</i>	5.2	4.8	5.1	4.4	3.4	2.9	2.4	1.9
AIDS Cases Cases	120	107	124	120	86	89	89	39
(DTP Reports) <i>Rate per 100,000</i>	2.9	2.5	2.9	2.7	1.9	2	1.9	0.8
AIDS Cases Cases	135	121	114	122	99	79	73	–
(BCCDC Reports) <i>Rate per 100,000</i>	3.2	2.9	2.6	2.8	2.2	1.7	1.6	–

Indicator 13: HIV-Related Mortality

	2004	2005	2006	2007	2008	2009	2010	2011
British Columbia Overall	105	146	142	100	79	63	54	59
Per 100 HIV+ Population	1.03	1.4	1.34	0.93	0.72	0.56	0.47	0.5
Per 100,000 Population	2.5	3.43	3.29	2.28	1.8	1.41	1.19	1.29