



BRITISH COLUMBIA  
CENTRE *for* EXCELLENCE  
*in* HIV/AIDS

# HIV MONITORING QUARTERLY REPORT **FOR FRASER HEALTH**

FOURTH QUARTER 2016



BC Centre for Disease Control  
An agency of the Provincial Health Services Authority



First Nations Health Authority  
Health through wellness



fraserhealth



Interior Health  
Every person matters



island health



northern health  
the northern way of caring



How you want to be treated.



Vancouver  
CoastalHealth  
Promoting wellness. Ensuring care.



## Foreword

As part of the BC Centre for Excellence (BC-CFE) in HIV/AIDS's mandate to evaluate the outcomes of STOP HIV/AIDS programming in BC, we have developed quarterly HIV/AIDS monitoring reports. These reports provide up-to-date data on a variety of key HIV-related surveillance and treatment indicators. Selection of these indicators was achieved through a collaborative process with various Health Authority (HA) representatives. There are six reports in total, one for each HA and one for the province of BC as a whole. In addition, there is a technical report which explains how each HIV indicator is calculated. Data used in these reports come from the British Columbia Centre for Disease Control (BCCDC), MSP billings, hospitalization data from the Discharge Abstract Database, the Sunquest Laboratory database at the Provincial Public Health Microbiology and Reference Laboratory, Providence Health Care laboratory, BC Vital Statistics, and the BC-CFE Drug Treatment Program (DTP) Database.

The objectives of these reports are to:

1. Provide timely HA-specific information on key HIV indicators which will guide and inform HIV leaders and innovators in the development of future HIV interventions and programs which will ultimately lead to decreasing the burden of HIV in BC. The indicators will reflect ongoing or past successful public health interventions and highlight areas in the HIV care spectrum which require further attention and support.
2. Highlight limitations in our current data due to incomplete or time lagged data and to develop future strategies to improve complete and timely data capture.

These reports are produced for the benefit of individual HA's. As such, we are enthusiastic about your involvement and cooperation regarding the development of these monitoring reports. Please forward your comments and queries to Irene Day, Director of Operations at the BC-CFE at [iday@cfenet.ubc.ca](mailto:iday@cfenet.ubc.ca).

## List of Indicators

Indicator 1. HIV Testing Episodes

Indicator 2. HIV Testing Rate

Indicator 3. New HIV Diagnoses

Indicator 4. Stage of HIV Infection at Diagnosis

Indicator 5. HIV Cascade of Care

Indicator 6. Programmatic Compliance Score (PCS)

Indicator 7. New Antiretroviral Therapy Starts

Indicator 8. CD4 Cell Count at ART Initiation

Indicator 9. Active and Inactive Drug Treatment Program (DTP) Participants

Indicator 10. Antiretroviral Adherence

Indicator 11. Resistance Testing and Results

Indicator 12. AIDS-Defining Illness

Indicator 13. HIV-Related Mortality

# Table of Contents

## Acknowledgements and Contributions

### BC Provincial STOP Program:

#### A Note on Monitoring and Interpreting Hiv Indicators

**Indicator 1      Hiv Testing Episodes** *All HIV Testing Episodes reflect non-prenatal tests. All prenatal tests have been removed.*

Figure 1.1      Hiv Test Episodes for Fraser Health, 2012 Q1–2016 Q4

Figure 1.2      Hiv Test Episodes for Fraser Health by Gender,  
2012 Q1–2016 Q4

Figure 1.3      Hiv Test Episodes for Fraser Health by Age Category, 2012 Q1–2016 Q4

Figure 1.4      Point-of-Care Hiv Tests for Fraser Health, 2012 Q1–2016 Q4

Figure 1.5      Hiv Test Episodes by HSDA for Fraser Health, 2012 Q1–2016 Q4

Figure 1.6      Hiv Test Episodes for Non-Prenatal Females in Fraser Health by HSDA, 2012 Q1–2016 Q4

Figure 1.7      Hiv Test Episodes for Males in Fraser Health by HSDA, 2012 Q1–2016 Q4

**Indicator 2      Hiv Testing Rates** *All HIV Testing Rates reflect non-prenatal tests. All prenatal tests have been removed.*

Figure 2.1      Rate of Hiv Testing for Fraser Health and HSDA's, 2009–2015

Figure 2.2      Rate of Hiv Testing for Fraser Health by Gender, 2009–2015

Figure 2.3      Rate of Hiv Testing for Fraser Health by Age Category, 2009–2015

**Indicator 3      New Hiv Diagnoses**

Figure 3.1      New Hiv Diagnoses for Fraser Health, 2012 Q1–2016 Q4

Figure 3.2      New Hiv Diagnoses for Fraser Health by Gender, 2012 Q1–2016 Q4

Figure 3.3      New Hiv Diagnoses for Fraser Health by Age Category, 2012 Q1–2016 Q4

Figure 3.4      New Hiv Diagnoses for Fraser Health by Exposure Category, 2012 Q1–2016 Q2

Figure 3.5      New Hiv Diagnoses for Fraser Health by HSDA, 2012 Q1–2016 Q4

**Indicator 4      Stage of Hiv Infection at Diagnosis** *Stage definitions have been altered to remove AIDS diagnosis data. Individuals previously classified as Stage 3 have been re-classified based on CD4 cell count.*

Table 1      Staging Classifications of Infection at Time of Hiv Diagnosis Based on CDC Hiv  
Surveillance Case Definitions

Figure 4.1      Stage of Hiv Infection at Diagnosis for Fraser Health, 2011–2015

Figure 4.2      Stage of Hiv Infection at Diagnosis for Fraser Health by Gender, 2011–2015

Figure 4.3      Stage of Hiv Infection at Diagnosis for Fraser Health by Age Category, 2011–2015

Figure 4.4      Stage of Hiv Infection at Diagnosis for Fraser Health by Exposure Category,  
2011–2015

**Indicator 5      Hiv Cascade of Care**

Figure 5.1      Estimated Cascade of Care for Fraser Health, Year Ending 2016 Q4

Figure 5.2      Estimated Cascade of Care for Fraser Health by Gender, Year Ending 2016 Q4

Figure 5.3      Estimated Cascade of Care for Fraser Health by Age Category, Year Ending 2016 Q4

Figure 5.4	Estimated Cascade of Care for Fraser Health by MSM Status, Year Ending 2016 Q4
Figure 5.5	Estimated Cascade of Care for Fraser Health by Age Category and MSM Status, Year Ending 2016 Q4
Figure 5.6	Estimated Cascade of Care for Fraser Health by PWID Status, Year Ending 2016 Q4
Figure 5.7	Estimated Cascade of Care for Fraser Health by HSDA, Year Ending 2016 Q4
<b>Indicator 6</b>	<b>Programmatic Compliance Score (PCS)</b>
Table 2	Probability of Mortality, Immunologic Failure and Virologic Failure Based on the Programmatic Compliance Score
Figure 6.1	PCS Components for Fraser Health, 2015 Q1–2016 Q4 <ul style="list-style-type: none"> <li>Less than 3 CD4 Tests in First Year</li> <li>Less than 3 Viral Load Tests in First Year</li> <li>Not Having Drug Resistance Testing at Baseline</li> <li>Non-Recommended Antiretroviral Therapy Regimen (ART)</li> <li>Baseline CD4 &lt; 200 cells/μL</li> <li>Not Achieving Viral Suppression at 9 Months</li> </ul>
Figure 6.2	Historical Trends for Pcs Score for Fraser Health, 2015 Q1–2016 Q4
<b>Indicator 7</b>	<b>New Antiretroviral Therapy Starts in Fraser Health</b>
Figure 7	BC-CfE Drug Treatment Program Enrollment: New Antiretroviral Participants for Fraser Health, 2015 Q1–2016 Q4
<b>Indicator 8</b>	<b>CD4 Cell Count at ART Initiation</b>
Figure 8	CD4 Cell Count at ART Initiation for Fraser Health, 2015 Q1–2016 Q4
<b>Indicator 9</b>	<b>Active and Inactive Drug Treatment Program (DTP) Participants</b>
Table 3	Distribution of People on ART in Fraser Health, 2016 Q4
Figure 9	Active and Inactive DTP Participants for Fraser Health, 2015 Q1–2016 Q4
<b>Indicator 10</b>	<b>Antiretroviral Adherence</b>
Figure 10	Distribution of Individuals by Adherence Level in 1st Year of Therapy, Based on Pharmacy Refill Compliance for Fraser Health, 2015 Q1–2016 Q4
<b>Indicator 11</b>	<b>Resistance Testing and Results</b>
Figure 11	Cumulative Resistance Testing Results by Resistance Category for Fraser Health, 2015 Q1–2016 Q4
<b>Indicator 12</b>	<b>AIDS-Defining Illness</b>
Figure 12	AIDS Case Rate and Reports for Fraser Health, 2008–2015
<b>Indicator 13</b>	<b>HIV-Related Mortality</b>
Figure 13	HIV-Related Deaths by Year for Fraser Health, 2004–2011

# Acknowledgements and Contributions



BRITISH COLUMBIA  
CENTRE *for* EXCELLENCE  
*in* HIV/AIDS

**British Columbia Centre for Excellence in HIV/AIDS (BC-CFE):** The BC-CFE is responsible for the conception, preparation and ongoing review of this quarterly report. The BC-CFE provides the data and outputs for Indicators 5 (Hiv Cascade of Care), 6 (Programmatic Compliance Score), 7 (New Antiretroviral Starts), 8 (CD4 Cell Count at ART Initiation), 9 (Active and Inactive Drug Treatment Program Participants), 10 (Antiretroviral Adherence Level), 11 (Resistance Testing Results by Resistance Category), 12 (AIDS-Defining Illness), and 13 (HIV-Related Mortality). The BC-CFE database provides PVL and CD4 cell count testing data, as well as ART use. All PVL measurements in BC are performed at the St Paul's Hospital virology laboratory, thus PVL data capture is 100%. An estimated 80% of all CD4 count measurements performed in the province are captured in the BC-CFE data holdings. The STOP HIV/AIDS Technical Monitoring Committee–BC-CFE is responsible for oversight of the monitoring report. James Nakagawa is responsible for compiling and publishing this report. Lilith Swetland is the editor of this report. Paul Sereda, Dr. Viviane Lima and Nada Gataric perform analysis of Indicators 5–13. This report was conceived and guided by Dr. Julio Montaner.



BC Centre for Disease Control  
An agency of the Provincial Health Services Authority

**British Columbia Centre for Disease Control (BCCDC):** The BCCDC provides the data and outputs for Indicator 1 (Hiv Testing Episodes), Indicator 2 (Hiv Testing Rate), Indicator 3 (New Hiv Diagnoses), Indicator 4 (Stage of Hiv at Diagnosis) and Indicator 12 (AIDS-Defining Illness). The BCCDC is the single provincial agency that centralizes all HIV surveillance through the Public Health Microbiology and Reference Laboratory, which does more than 90% of all HIV screening tests in BC and all confirmatory testing. Olga Mazo, Theodora Consolacion and Dr. Jason Wong are responsible for outputs for Indicators 1–4.

## Other Data Sources:

The above databases were supplemented with:

- (I) The BC Vital Statistics database which was used to calculate Indicator 5. The Hiv Cascade of Care and Indicator 13. Hiv-Related Mortality.
- (II) Linkage and preparation of the de-identified individual-level database used for calculating Indicator 5. The Hiv Cascade of Care was facilitated by the British Columbia Ministry of Health.
- (III) The Statistics Canada database: BC and HIV-positive population counts were acquired through the statistics Canada website to calculate HIV-specific mortality rates for Indicator 13. Hiv-Related Mortality.

# Membership of the STOP HIV/AIDS Technical Monitoring Committee–BC-CfE

Dr. Rolando Barrios, *Chair*, BC-CfE

Dr. Kate Heath, BC-CfE

Dr. Bohdan Nosyk, BC-CfE

Dr. Viviane Dias Lima, BC-CfE

Irene Day, BC-CfE

Dr. Jean Shoveller, BC-CfE

Dr. Jason Wong, BCCDC

Dr. Mel Krajden, BCCDC

Salman Klar, FHA

Gillian Frosst, IHA

Kari Harder, NHA

Dr. Neora Pick, PHSA

Dr. Reka Gustafson, VCHA

Dr. Melanie Rusch, VIHA

Robert Parker, FNHA



# The Seek and Treat for Optimal Prevention (STOP) HIV/AIDS BC Provincial Program: A Note on Monitoring and Interpreting HIV Indicators

The Seek and Treat for Optimal Prevention (STOP) of HIV/AIDS programme is a provincial initiative to improve HIV diagnosis and care delivery in BC through increased HIV-specific funding to all Health Service Delivery Areas (HSDA's) across BC. The STOP provincial programme is an expansion of a four-year STOP pilot project which was implemented in two Health Service Delivery Areas in March 2010; the Vancouver HSDA which bears the largest burden of the HIV epidemic in the province and the Northern Interior HSDA which bears a high burden of HIV-related mortality. The STOP pilot project demonstrated the urgent need for improved efforts in early diagnosis of HIV and timely initiation of antiretroviral therapy (ART) initiation.

The expansion to a province-wide programme was announced on November 30th, 2013 by the BC Ministry of Health with roll out of funding beginning on April 1st, 2013. This funding is intended to be used in the implementation and evaluation of HIV-related diagnosis and care initiatives within individual HA's. Goals of the project include: 1. A reduction in the number of new HIV infections in BC; 2. Improvements in the quality, effectiveness, and reach of HIV prevention services; 3. An increase in early diagnosis of HIV; 4. A reduction in AIDS cases and HIV-related mortality.

The goals of HA-led STOP-funded initiatives are to work toward achieving these goals. To these ends some outcome measures or indicators of progress have been drafted that should be considered in the design and implementation phases of these initiatives.

# HIV Testing Episodes and Rates

In this section, the number of HIV test episodes and point of care (POC) HIV tests conducted each quarter in BC is shown. In general terms the goal is to increase the number of tests performed and to maximize testing efficiency. Test episodes are allocated by region according to where the test is performed.

## Indicator 1. HIV Testing Episodes

Figure 1.1 HIV Test Episodes for Fraser Health




Figure 1.2 HIV Test Episodes by Gender for Fraser Health <sup>1</sup>




Figure 1.3 HIV Test Episodes by Age Category for Fraser Health <sup>1,2</sup>





Figure 1.4 Point-of-Care HIV Tests for Fraser Health



1 Data Source: The BC Public Health Microbiology and Reference Laboratory (BCPHMRL) courtesy of the BC Centre for Disease Control (BCCDC).

Limitation: Repeat tests in individuals who test using various identifiers may not be identified and these individuals may be counted more than once.

2 Testing does not include point of care tests.

Figure 1.5 HIV Test Episodes for Fraser Health by HSDA <sup>1</sup>




Figure 1.6 HIV Test Episodes for Non-prenatal Females in Fraser Health by HSDA <sup>1</sup>





Figure 1.7 HIV Test Episodes for Males in Fraser Health by HSDA <sup>1</sup>



## Indicator 2. HIV Testing Rates

Figure 2.1 Rate of HIV Testing for Fraser Health and HSDAs <sup>2</sup>




Figure 2.2 Rate of HIV Testing by Gender for Fraser Health <sup>2</sup>





Figure 2.3 Rate of HIV Testing by Age Category for Fraser Health<sup>2</sup>



<sup>2</sup> Testing does not include point of care tests.

# New HIV Diagnoses

Trends in HIV diagnoses by gender and exposure category are described. Interpreting HIV diagnoses must be done with consideration that trends are influenced by both changes in testing rate as well as changes in transmission rates. It is important to note that new HIV diagnoses cases and rates are not synonymous with HIV incidence as a person may have become infected with HIV long before they tested positive for HIV. However, as there is no reliable method for measuring HIV incidence, we follow trends in HIV diagnoses.

## Indicator 3. New HIV Diagnoses

Figure 3.1 New HIV Diagnoses for Fraser Health<sup>3</sup>





Figure 3.2 New HIV Diagnoses for Fraser Health by Gender<sup>3</sup>



<sup>3</sup> Data Source: BCCDC. When present, "By Provider Address" is graphed as dashed line in same colour.



Figure 3.3 New HIV Diagnoses for Fraser Health by Age Category<sup>3</sup>




Figure 3.4 New HIV Diagnoses for Fraser Health by Exposure Category<sup>3,4</sup>





Figure 3.5 New HIV Diagnoses for Fraser Health by HSDA<sup>3</sup>



<sup>3</sup> Data Source: BCCDC. When present, "By Provider Address" is graphed as dashed line in same colour.

<sup>4</sup> MSM=men who have sex with men; PWID=people who inject drugs; HET=heterosexual. NIR=No identified risk/exposure.

## Stage of HIV Infection at Diagnosis

Classification of stage of HIV infection, in the absence of information regarding recent testing history, is reliant on clinical information available at the time of diagnosis, including first CD4+ cell count and laboratory results suggestive of acute HIV infection (Table 1). The benefits of Treatment as Prevention (TasP) are maximized when antiretroviral therapy (ART) is initiated at high CD4 cell counts. Accordingly, it is preferable that individuals newly diagnosed with HIV be in the early stages of HIV infection (stage 0 or 1) to allow for early ART initiation.

*N.B. Interpretation of Stage of HIV Infection at Diagnosis should proceed with caution. Early increases in diagnosis at late stage (i.e., low CD4 counts) may represent a “catching up” of previously missed long term infected individuals rather than a trend toward diagnosis at later stage of infection.*

## Indicator 4. Stage of HIV Infection at Diagnosis

Table 1: Staging Classifications of Infection at Time of HIV Diagnosis Based on CDC HIV Surveillance Case Definitions

Stage	Criteria	
0	Laboratory criteria met for acute HIV infection, or previous negative or indeterminate HIV test within 180 days of first confirmed positive HIV test.	
1	Stage 0 not met <i>and</i>	CD4 ≥500
2a		CD4 350–499
2b		CD4 200–349
3		CD4 <200
Unknown		No available CD4

Figure 4.1 Stage of HIV Infection at Diagnosis for Fraser Health <sup>5</sup>





Figure 4.2 Stage of HIV Infection at Diagnosis by Gender for Fraser Health <sup>5</sup>



	Fraser Health					Female					Male				
	2011	'12	'13	'14	'15	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15
Stage 0	10	3	6	6	6	1	0	0	0	1	9	3	6	6	5
Stage 1	8	6	10	8	15	2	1	4	3	3	6	5	6	5	12
Stage 2a	3	5	6	8	6	1	3	1	2	0	2	2	5	6	6
Stage 2b	11	6	8	2	6	3	1	1	1	2	8	5	7	1	4
Stage 3	16	9	15	12	13	2	2	1	4	1	14	7	14	8	12
Unknown	1	2	2	10	6	1	0	0	4	1	0	2	2	6	5
Total (n=)	49	31	47	46	52	10	7	7	14	8	39	24	40	32	44

Figure 4.3 Stage of HIV Infection at Diagnosis by Age Category for Fraser Health <sup>5</sup>





Figure 4.4 Stage of HIV Infection at Diagnosis by Exposure Category for Fraser Health <sup>5,6</sup>



	< 30 years					30-39 years					40-49 years					≥ 50 years					MSM					PWID					Heterosexual					Other					NIR/Unknown						
	2011	12	13	14	15	11	12	13	14	15	11	12	13	14	15	11	12	13	14	15	11	12	13	14	15	11	12	13	14	15	11	12	13	14	15	11	12	13	14	15							
Stage 0	4	0	4	4	1	2	1	0	1	1	2	1	1	0	1	2	1	1	1	3	6	2	6	6	3	1	0	0	0	0	3	1	0	0	0	0	0	0	0	0	0	0	0	0			
Stage 1	1	1	3	3	9	5	0	3	3	2	1	4	1	1	1	1	1	3	1	3	3	2	3	3	3	11	1	0	2	2	0	4	3	5	3	4	0	1	0	0	0	0	0	0	0	0	
Stage 2a	0	0	2	1	1	2	1	3	3	1	1	1	1	1	2	0	3	0	3	2	2	2	3	4	0	0	1	1	1	0	0	3	1	4	1	1	0	2	0	0	0	0	0	0	0		
Stage 2b	1	2	3	0	1	2	1	0	0	2	3	1	2	0	2	5	2	3	2	1	4	2	5	0	2	1	1	0	0	1	6	3	3	2	3	0	0	0	0	0	0	0	0	0	0		
Stage 3	0	0	1	1	0	4	3	3	3	2	8	3	6	4	4	4	3	5	4	7	5	2	5	5	3	2	3	3	1	0	8	4	5	6	9	1	0	1	0	0	0	0	1	0	1		
Unknown	1	1	0	2	0	0	0	0	2	2	0	1	2	5	4	0	0	0	1	0	0	1	0	4	3	0	0	0	2	0	1	0	2	4	3	0	0	0	0	0	0	0	0	1	0	0	0
Total (n=)	7	4	13	11	12	15	6	9	12	10	15	11	13	11	14	12	10	12	12	16	20	11	21	21	26	5	4	6	6	2	22	14	16	19	23	2	1	3	0	0	0	1	1	0	1		

<sup>5</sup> Data Source: BCCDC

<sup>6</sup> MSM=men who have sex with men; PWID=people who inject drugs; HET=heterosexual. NIR=No identified risk/exposure.

# HIV Cascade of Care

## Indicator 5. HIV Cascade of Care

The success of seek, test, treat and retain (STTR) strategies like STOP is reliant on early diagnosis of HIV, linking newly diagnosed HIV-positive persons with ongoing care, retaining persons in HIV-care; initiating ART based on best evidenced practices and maintaining optimal ART adherence to ensure a suppressed viral load. These stages of HIV-care can be summarized as: 1. HIV diagnosis, 2. Linked to HIV care, 3. Retained in HIV care, 4. On ART, 5. Adherent to ART and 6. Achieving a suppressed VL; collectively, they are referred to as the cascade of care. Attrition between any of these stages of HIV-care means a reduction in the potential of ART as a benefit to the HIV-positive individual and as an HIV transmission prevention method on a population level. Thus, when interpreting trends in the cascade of care, we strive to see increases along each step of the cascade of care (i.e. reduced attrition) with the ultimate goal being 100% within each stage of the cascade. Monitoring the Cascade of Care provides a picture as to where deficiencies lie in the delivery and uptake of HIV-care. In this section we present the cascade of care for the period 2015 Q4–2016 Q3 in Fraser Health and stratified by sex and age.

Figure 5.1 Estimated Cascade of Care for Fraser Health, Year Ending 2016 Q4 <sup>7</sup>





Figure 5.2 Estimated Cascade of Care for Fraser Health by Gender, Year Ending 2016 Q4 <sup>7</sup>



<sup>7</sup> Data is for the period 2015 Q4–2016 Q3.


Data Sources:

- i British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- ii Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider. If the most recent HA of residence is not updated then the designated HA may be incorrect.


NB: Transgender have been assigned to their biological sex.

Figure 5.3 Estimated Cascade of Care for Fraser Health by Age Category, Year Ending 2016 Q4 <sup>8</sup>



<sup>8</sup> Data is for the period 2015 Q4–2016 Q3.  
Data Sources:  
i British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).  
ii Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).  
Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider.  
If the most recent HA of residence is not updated then the designated HA may be incorrect.

Figure 5.4 Estimated Cascade of Care for Fraser Health by MSM Status, Year Ending 2016 Q4 <sup>9</sup>




<sup>9</sup> Data is for the period 2015 Q4–2016 Q3.

Data Sources:

- i British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- ii Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider. If the most recent HA of residence is not updated then the designated HA may be incorrect.

Figure 5.5 Estimated Cascade of Care for Fraser Health by Age Category and MSM Status, Year Ending 2016 Q4 <sup>9</sup>




<sup>9</sup> Data is for the period 2015 Q4–2016 Q3.

Data Sources:

- i British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- ii Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider. If the most recent HA of residence is not updated then the designated HA may be incorrect.


Figure 5.6 Estimated Cascade of Care for Fraser Health by PWID Status, Year Ending 2016 Q4 <sup>9</sup>



<sup>9</sup> Data is for the period 2015 Q4–2016 Q3.  
 Data Sources:  
 i British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).  
 ii Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).  
 Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider.  
 If the most recent HA of residence is not updated then the designated HA may be incorrect.



Figure 5.7 Estimated Cascade of Care for Fraser Health by HSDA, Year Ending 2016 Q4 <sup>9</sup>



<sup>9</sup> Data is for the period 2015 Q4–2016 Q3.

Data Sources:

- i British Columbia Centre for Excellence Drug Treatment Program (DTP) Database (ARV use, VL and CD4 count).
- ii Administrative data (ex. MSP billings; hospitalization data from the Discharge Abstract Database (DAD)).

Limitations: HA assignment is based on the most recent HA of residence of the patient, if not available of the HIV-care provider. If the most recent HA of residence is not updated then the designated HA may be incorrect.

# Programmatic Compliance Score

## Indicator 6. Programmatic Compliance Score (PCS)

The Programmatic Compliance Score (PCS) is a summary measure of risk of future death, immunologic failure and virologic failure from all causes for people who are starting ART for the first time. It is composed of patient- and physician-driven effects. PCS scores range from 0–6 with higher scores indicative of poorer health outcomes and greater risk of death. Table 2 provides mortality, immunologic failure and virologic failure probabilities for given PCS scores. We interpret an individual with a  $PCS \geq 4$  as being 22 times more likely to die, almost 10 times more likely to have immunologic failure and nearly 4 times as likely to demonstrate virologic failure compared to those individuals with a PCS score of 0. A detailed description of how the PCS score is calculated and its validation can be found in the technical report. In short, PCS scores are calculated by summing the results (yes=1, no=0) of six un-weighted non-performance indicators based on IAS–USA treatment guidelines:

1. having <3 CD4 cell count tests in the first year after starting antiretroviral therapy (ART);
2. having <3 plasma viral load (VL) tests in the first year after starting ART;
3. not having drug resistance testing done prior to starting ART;
4. starting on a non-recommended ART regimen;
5. starting therapy with  $CD4 < 200$  cells/ $\mu$ L; and
6. not achieving viral suppression within 9 months since ART initiation.

In this section we provide PCS scores and their components over time for the province of BC. A decline to 0%, (i.e., **all individuals having a score of 0**) is the eventual goal.

Table 2. Probability of Mortality, Immunologic Failure and Virologic Failure based on the Programmatic Compliance Score

Programmatic Compliance Score	Mortality Risk Ratio (95% Confidence Interval)	Immunologic Failure Risk Ratio (95% CI)	Virologic Failure Risk Ratio (95% CI)
0 (Best score)	1 (–)	1 (–)	1 (–)
1	3.81 (1.73–8.42)	1.39 (1.04–1.85)	1.32 (1.05–1.67)
2	7.97 (3.70–17.18)	2.17 (1.54–3.04)	1.86 (1.46–2.38)
3	11.51 (5.28–25.08)	2.93 (1.89–4.54)	2.98 (2.16–4.11)
4 or more (Worst score)	22.37 (10.46–47.84)	9.71 (5.72–16.47)	3.80 (2.52–5.73)

Reference: Lima VD, Le A, Nosyk B, Barrios R, Yip B, et al. (2012) Development and Validation of a Composite Programmatic Assessment Tool for HIV Therapy. PLoS ONE 7(11): e47859. doi:10.1371/journal.pone.0047859

Figure 6.1 PCS Components for Fraser Health, 2015 Q1–2016 Q4 <sup>10</sup>





Figure 6.2 Historical Trends for PCS Score for Fraser Health, 2015 Q1–2016 Q4 <sup>10,11</sup>



<sup>10</sup> Data Source: British Columbia Centre for Excellence Drug Treatment Program (DTP) Database. Limitations: CD4 cell count capture is approximately 80%.


<sup>11</sup> Each quarter's data is calculated as the sum of the 4 quarters leading up to it. e.g. 2013 Q1 is calculated from 2012 Q2 – 2013 Q1. NB: A score of 0 is the best score and a score of 4 or more is the worst score.

# Antiretroviral Uptake

In this section we present trends in ART uptake, the number and proportion of new HIV treatment initiations and the number of active and inactive DTP participants. Trends in ART uptake should be interpreted under the consideration of changing BC HIV treatment guidelines. BC HIV treatment guidelines are updated regularly by the BC-CfE Therapeutic Guidelines Committee and reflect those of the International AIDS Society. Most recent changes were made in 2012 and HIV treatment is now recommended for all HIV-positive adults regardless of CD4 cell count; as evidence demonstrates that early initiation of HIV treatment maximizes both the individual's health outcomes as well as the potential of ART as a form of HIV transmission prevention at a population level. As such, trends in the number and proportion of persons on ART and new ART starts (in both naïve and experienced persons) are expected to increase over time at higher CD4 cell counts.


## Indicator 7. New Antiretroviral Therapy Starts in Fraser Health

Figure 7 BC-CfE Drug Treatment Program Enrollment: New ART Participants in Fraser Health, 2015 Q1–2016 Q4 <sup>12</sup>



## Indicator 8. CD4 Cell Count at ART Initiation

Figure 8 CD4 Cell Count at ART Initiation of ART-Naïve DTP Participants in Fraser Health, 2015 Q1–2016 Q4 <sup>13</sup>



<sup>12</sup> Data Source: Drug Treatment Program Database  
Limitation: DTP participants are designated to an HA based on most current residence provided by the participant.


<sup>13</sup> Data Source: Drug Treatment Program Database  
Limitations: CD4 cell count data is approximately 80% complete.

## Indicator 9. Active and Inactive DTP Participants

Table 3. Distribution of People on ART for Fraser Health, 2016 Q4 <sup>14</sup>

Age	< 30	81
	30–39	272
	40–49	513
	≥ 50	953
Gender	Male	1407
	Female	412
Exposure	MSM	598
	PWID	453
Total		1819

Figure 9 Active and Inactive DTP Participants in Fraser Health, 2015 Q1–2016 Q4 <sup>15</sup>



<sup>14</sup> Data Source: Drug Treatment Program Database

Limitation: DTP participants are designated to an HA based on most current residence provided by the participant.

Definition:

'On antiretroviral therapy' defined as being on treatment in the current quarter

<sup>15</sup> Active DTP participants: An individual who has had medication prescribed at least once in the preceding quarter.


Inactive DTP participants: Persons no longer prescribed drugs through the HIV/AIDS Drug Treatment Program in the last quarter.

# Antiretroviral Adherence Level

In this section we present trends in prescription refill adherence levels for individuals in their first year of treatment. Given that the benefits of ART are compromised in the presence of imperfect ART adherence, we expect to see the proportion of persons on ART achieving **near perfect adherence** (ie.  $\geq 95\%$ ) to increase with time. Furthermore, it is important that trends in the proportion of ART users achieving prescription refill adherence of  $\geq 95\%$  keep pace with new ART starts and increase among those continuing on ART.

## Indicator 10. Antiretroviral Adherence

Figure 10 Distribution of Individuals by Adherence Level in 1st Year of Therapy, Based on Pharmacy Refill Compliance for Fraser Health, 2015 Q1–2016 Q4 <sup>16</sup>



<sup>16</sup> Data Source: Drug Treatment Program Database


Limitation: Prescription refill adherence is used as a proxy for patient adherence.

# Resistance Testing and Results

## Indicator 11. Resistance Testing and Results

In this section, we present trends in cumulative resistance testing by resistance category: **Suppressed** (where a DTP participant's viral load is too low to be genotyped); **Wild Type** (where no HIV treatment resistances were discovered), **Never Genotyped**, and Resistances to **one**, **two**, **three**, or **four** HIV treatment classes. Resistance testing prior to ART initiation is recommended in the BC HIV treatment primary care guidelines. Thus, it is expected that trends over time should find all persons enrolled in the DTP to have been genotyped. Trends over time should also show an increase in the proportion of DTP participants achieving a suppressed status and an increase in resistance testing should not lead to an increase in the number of ART resistances occurring.

Figure 11 Cumulative Resistance Testing Results by Resistance Category for Fraser Health, 2015 Q1–2016 Q4 <sup>17</sup>



<sup>17</sup> Data Source: Drug Treatment Program Database


Limitation: DTP participants are designated to a HA based on most current residence provided by the participant.

# AIDS-Defining Illness

## Indicator 12. AIDS-Defining Illness

Improvements in ART and the expansion of ART province-wide has led to very low numbers of recorded AIDS cases across BC. However, interpreting trends in AIDS cases is challenging as AIDS reporting is passive in BC and it is likely that they are under-reported across all Health Authorities. In addition to under-reporting, methods of reporting AIDS cases are inconsistent across HA's and do not truly reflect the current reality of new AIDS diagnoses. Efforts will need to be made to improve under- and inconsistent reporting of AIDS cases across all HA's. The table below shows AIDS cases using three definitions. First, AIDS cases were defined as the number of physician-reported AIDS defining illness (ADI) in a given year. AIDS case reporting is a passive process and physicians can voluntarily report AIDS cases to the BCCDC or DTP. As such, we have plotted both **BCCDC reports** and **DTP reported AIDS cases**. We also show the proportion of persons **initiating ART with a CD4<200 cells/μL**.

Figure 12 AIDS Case Rate and Reports for Fraser Health <sup>18</sup>



<sup>18</sup> Data Source: DTP AIDS cases are obtained from the Drug Treatment Program Database; BCCDC AIDS cases are obtained from the BC-CDC; CD4<200 at ART initiation data came from the DTP database. Indicator 12 also reflects information from BC Vital Statistics. As this information is made available to BC-CFE, we use it to inform the development and refinement of this indicator.

*Limitation:* AIDS case reporting was investigated using 3 definitions: First, using AIDS cases reported in AIDS case report forms from the DTP; Second, using AIDS cases reported via the BCCDC and third, using a CD4 cell count of <200 cells/μL at time of ART initiation using DTP data. AIDS case reporting is passive in BC, thus; AIDS case reporting is not well captured. The DTP sends out AIDS reporting forms to physicians annually. The BCCDC uses DTP AIDS case reports as well as physician AIDS case reports made directly to the BCCDC. Interpreting AIDS case reports should be done with these limitations in mind. AIDS data is updated annually as very few AIDS cases reports are reported in general and trends would be difficult to notice if reported quarterly.



## HIV-Related Mortality

### Indicator 13. HIV-Related Mortality

This indicator is currently under revision.

## APPENDICES

Indicator 1: Test Episodes (thousands)		2012				2013				2014				2015				2016			
		Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
Fraser Health		10.1	10.1	10.8	11.6	13.5	14.0	13.7	12.9	13.6	14.5	14.9	15.1	16.1	15.8	16.8	17.1	18.5	19.2	17.9	17.1
Gender	Female	4.7	4.9	5.3	5.7	6.6	6.8	6.8	6.1	6.5	6.9	7.2	7.4	7.6	7.6	8.1	8.1	8.8	9.3	8.6	8.2
	Male	5.3	5.1	5.4	5.7	6.6	7.0	6.7	6.5	6.9	7.1	7.3	7.5	8.1	7.8	8.4	8.6	9.3	9.6	9.1	8.7
Age	Other	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	< 30	3.6	3.6	3.9	3.8	4.1	4.3	4.4	4.1	4.0	4.3	4.5	4.4	4.6	4.6	5.1	5.1	5.1	5.5	5.4	5.2
	30–39	2.6	2.5	2.6	2.7	3.3	3.3	3.3	3.2	3.4	3.4	3.6	3.6	3.9	3.8	4.0	3.9	4.5	4.5	4.2	4.1
	40–49	1.8	1.8	1.8	2.1	2.4	2.5	2.4	2.2	2.4	2.5	2.5	2.5	2.8	2.6	2.8	2.8	3.1	3.1	2.8	2.8
	≥ 50	2.0	2.1	2.4	2.8	3.4	3.7	3.4	3.1	3.5	3.8	4.0	4.3	4.4	4.7	4.9	5.4	5.7	5.3	4.9	
POC Tests ( <i>not in thousands</i> )		54	121	31	158	296	187	182	302	254	426	377	253	423	383	288	333	347	284	200	176
Fraser East		1.5	1.4	1.5	1.5	1.6	1.7	1.7	1.6	1.8	1.7	1.9	2.5	2.3	2.3	2.4	2.3	2.4	2.4	2.3	2.3
	Female	0.7	0.7	0.8	0.8	0.8	0.9	0.8	0.8	0.9	0.9	0.9	1.3	1.2	1.2	1.2	1.2	1.2	1.1	1.1	
	Male	0.7	0.7	0.7	0.7	0.7	0.8	0.8	0.8	0.9	0.8	0.9	1.2	1.1	1.1	1.2	1.1	1.2	1.2	1.1	
Fraser North		5.1	5.1	5.2	5.7	6.9	7.1	6.8	6.5	6.8	7.0	7.0	6.9	7.4	7.4	7.8	8.0	8.8	9.2	8.6	8.1
	Female	2.4	2.5	2.5	2.8	3.3	3.4	3.3	3.1	3.2	3.5	3.4	3.3	3.4	3.5	3.7	3.7	3.8	4.2	4.5	3.9
	Male	2.7	2.5	2.6	2.8	3.4	3.6	3.4	3.3	3.5	3.4	3.5	3.5	3.8	3.7	3.9	4.1	4.5	4.6	4.3	4.2
Fraser South		3.5	3.7	4.1	4.5	5.0	5.2	5.3	4.8	5.1	5.7	6.0	5.6	6.4	6.1	6.6	6.8	7.3	7.5	7.0	6.7
	Female	1.6	1.8	2.0	2.2	2.4	2.5	2.6	2.2	2.4	2.6	2.9	2.7	2.9	2.9	3.2	3.1	3.4	3.6	3.3	3.2
	Male	1.9	1.8	2.0	2.2	2.4	2.6	2.5	2.3	2.6	2.8	2.9	2.8	3.2	3.0	3.3	3.5	3.7	3.7	3.6	3.4

Indicator 2: <b>Rate of HIV Testing per 100,000</b>		2009	2010	2011	2012	2013	2014	2015
All Fraser Health		2251.1	2267.0	2330.7	2795.6	3316.2	3593.6	4014.6
Fraser East		2069.2	1997.6	1956.3	2249.2	2475.0	2947.6	3390.5
Fraser North		2707.1	2714.9	2809.4	3409.4	4293.9	4464.1	4851.4
Fraser South		1935.1	1993.3	2073.3	2489.6	2819.5	3113.9	3554.2
Gender	Female	2150.9	2171.0	2230.1	2768.4	3373.7	3614.2	4030.6
	Male	2342.9	2352.4	2422.6	2812.4	3248.4	3562.3	3987.0
Age	< 30	2215.9	2220.6	2244.2	2488.4	2689.3	2752.5	3060.6
	30–39	4370.1	4459.5	4509.5	5049.3	5666.9	6086.3	6611.5
	40–49	2658.3	2642.6	2717.3	3256.8	3896.4	4189.0	4702.8
	≥ 50	1145.0	1191.5	1320.7	1994.9	2797.9	3251.0	3709.1

Indicator 3: New HIV Diagnoses		2012				2013				2014				2015				2016			
		Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
Fraser Health	By Client Residence	11	10	10	14	8	14	23	19	15	11	16	17	18	17	15	12	19	20	19	15
	By Provider Address	10	5	7	8	9	8	14	15	11	10	12	13	11	11	15	14	17	15	14	14
Gender	Female	0	3	2	3	1	3	2	3	6	4	6	2	6	1	2	2	5	3	4	3
	Male	11	7	8	11	7	11	21	16	9	7	10	15	12	16	13	10	14	17	15	12
Age	< 30	3	1	1	2	1	3	5	6	5	1	4	4	3	4	3	4	6	3	8	2
	30–39	1	3	2	2	3	4	8	6	5	3	5	4	5	4	3	2	7	4	5	3
	40–49	3	2	5	4	2	4	6	4	3	3	1	4	2	6	3	3	4	4	3	5
	≥ 50	4	4	2	6	2	3	4	3	2	4	6	5	8	3	6	3	2	9	3	5
Exposure	MSM	6	5	6	5	5	8	10	11	5	3	8	11	7	8	8	6	7	9		
	PWID	4	0	0	1	1	1	2	5	3	1	0	3	0	0	1	0	0	0		
	HET	1	4	4	6	1	3	9	3	7	7	7	3	9	8	6	6	5	3		
	Other	0	0	0	1	0	1	2	0	0	0	1	0	0	0	0	0	2	0		
	NIR/Unknown	0	1	0	1	1	1	0	0	0	0	0	0	2	1	0	0	5	8		
Fraser East	By Client Residence	2	2	1	1	0	0	3	1	2	0	2	2	1	1	2	1	5	1	0	2
	By Provider Address	2	0	1	0	0	0	2	1	2	1	1	2	0	1	3	1	3	1	0	2
Fraser North	By Client Residence	3	6	5	6	6	9	14	9	8	5	5	10	9	10	5	5	6	7	9	8
	By Provider Address	3	4	4	2	7	4	8	9	7	4	4	9	7	4	5	6	5	4	7	7
Fraser South	By Client Residence	6	2	4	7	2	5	6	9	5	6	9	5	8	6	8	6	8	12	10	5
	By Provider Address	5	1	2	6	2	4	4	5	2	5	7	2	4	6	7	7	9	10	7	5

Indicator 4: **Stage of HIV Infection at Baseline**

	Fraser Health					Female					Male					< 30 years					30-39 years					40-49 years				
	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15
Stage 0	10	3	6	6	6	1	0	0	0	1	9	3	6	6	5	4	0	4	4	1	2	1	0	1	1	2	1	1	0	1
1	8	6	10	8	15	2	1	4	3	3	6	5	6	5	12	1	1	3	3	9	5	0	3	3	2	1	4	1	1	1
2a	3	5	6	8	6	1	3	1	2	0	2	2	5	6	6	0	0	2	1	1	2	1	3	3	1	1	1	1	1	2
2b	11	6	8	2	6	3	1	1	1	2	8	5	7	1	4	1	2	3	0	1	2	1	0	0	2	3	1	2	0	2
3	16	9	15	12	13	2	2	1	4	1	14	7	14	8	12	0	0	1	1	0	4	3	3	3	2	8	3	6	4	4
Unknown	1	2	2	10	6	1	0	0	4	1	0	2	2	6	5	1	1	0	2	0	0	0	0	2	2	0	1	2	5	4
Total	49	31	47	46	52	10	7	7	14	8	39	24	40	32	44	7	4	13	11	12	15	6	9	12	10	15	11	13	11	14

	≥ 50 years					MSM					Heterosexual					PWID					Other Exposure					NIR/Unknown				
	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15	'11	'12	'13	'14	'15
Stage 0	2	1	1	1	3	6	2	6	6	3	1	0	0	0	0	3	1	0	0	3	0	0	0	0	0	0	0	0	0	0
1	1	1	3	1	3	3	2	3	3	11	1	0	2	2	0	4	3	5	3	4	0	1	0	0	0	0	0	0	0	0
2a	0	3	0	3	2	2	2	2	3	4	0	0	1	1	1	0	3	1	4	1	1	0	2	0	0	0	0	0	0	0
2b	5	2	3	2	1	4	2	5	0	2	1	1	0	0	1	6	3	3	2	3	0	0	0	0	0	0	0	0	0	0
3	4	3	5	4	7	5	2	5	5	3	2	3	3	1	0	8	4	5	6	9	1	0	1	0	0	0	0	1	0	1
Unknown	0	0	0	1	0	0	1	0	4	3	0	0	0	2	0	1	0	2	4	3	0	0	0	0	0	0	1	0	0	0
Total	12	10	12	12	16	20	11	21	21	26	5	4	6	6	2	22	14	16	19	23	2	1	3	0	0	0	1	1	0	1

Indicator 5: <b>HIV Cascade of Care</b>			Diagnosed	Linked	Retained	On ARVs	Adherent	Suppressed
Fraser Health			2171	2085	1871	1785	1630	1409
Gender	Men		1650	1592	1435	1377	1284	1116
	Women		521	493	436	408	346	293
Age Category	< 30		121	97	85	77	64	51
	30-39		310	300	263	243	207	172
	40-49		610	588	519	496	450	371
	≥ 50		1130	1099	1005	969	909	815
MSM Status	MSM		655	642	609	583	549	495
	Non-MSM		792	768	689	653	581	480
	Unknown		724	674	573	549	500	434
Age Category and MSM Status	MSM	< 30	37	36	33	30	26	22
		30-39	93	90	80	73	62	56
		40-49	132	130	121	116	111	94
		≥ 50	393	386	376	364	350	323
	Non-MSM	< 30	10	10	9	7	5	5
		30-39	111	108	96	87	69	55
		40-49	271	261	235	223	195	149
		≥ 50	399	389	349	336	312	271
	Unknown	< 30	73	50	43	40	33	24
		30-39	106	102	88	83	76	61
		40-49	207	198	162	157	144	128
		≥ 50	338	324	280	269	247	221
PWID Status	PWID		532	520	478	451	391	305
	Non-PWID		1189	1160	1082	1038	975	869
	Unknown		450	405	311	296	264	235
HSDA	Fraser East		330	315	271	263	240	196
	Fraser North		954	903	808	765	702	607
	Fraser South		888	868	792	757	688	606

Indicator 6: <b>Programmatic Compliance Score (PCS)</b>	2015				2016			
	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
< 3 CD4 Tests	9.3%	7.0%	7.3%	9.5%	11.5%	11.2%	16.3%	21.7%
< 3 Viral Load Tests	3.5%	3.5%	4.9%	2.4%	9.2%	10.2%	10.9%	15.2%
No Baseline Genotype	3.5%	4.7%	3.7%	2.4%	5.7%	5.1%	5.4%	10.9%
Baseline CD4 < 200 cells/μL	27.9%	26.7%	19.5%	25.0%	21.8%	23.5%	26.1%	22.8%
Non-Recommended ART	1.2%	1.2%	0.0%	1.2%	3.4%	4.1%	5.4%	7.6%
Non Viral Suppression at 9 Mo.	25.6%	20.9%	23.2%	21.4%	32.2%	37.8%	38.0%	37.0%
PCS Score: 0	41	43	44	44	38	38	36	30
PCS Score: 1	32	33	31	31	34	42	33	36
PCS Score: 2	10	8	4	6	6	8	12	13
PCS Score: 3	3	2	3	3	9	8	7	8
PCS Score: 4 or more	0	0	0	0	0	2	4	5
<b>Total (n=)</b>	<b>86</b>	<b>86</b>	<b>82</b>	<b>84</b>	<b>87</b>	<b>98</b>	<b>92</b>	<b>92</b>

Indicator 7: **New DTP ARV Participants**

First Starts	31	27	17	23	20	23	30	13
Experienced Starts	37	35	38	36	31	36	29	40

Indicator 8: **CD4 Cell Count Initiation for ARV-Naïve DTP Participants**

CD4 ≥ 500	14	8	4	10	3	11	11	6
CD4 350-499	6	3	2	2	5	2	5	0
CD4 200-349	4	7	2	5	2	6	3	6
CD4 50-199	4	4	4	1	6	2	9	0
CD4 < 50	0	5	4	2	3	2	1	1
CD4 MED	490	310	180	500	330	479	375	349
<b>Total (n=)</b>	<b>28</b>	<b>27</b>	<b>16</b>	<b>20</b>	<b>19</b>	<b>23</b>	<b>29</b>	<b>13</b>

Indicator 9: **Active and Inactive DTP Participants**

Active DTP Participants	1706	1716	1748	1760	1763	1763	1788	1819
Inactive DTP Participants	250	252	241	247	256	260	274	258

Indicator 10: **Antiretroviral Adherence**

≥ 95%	23	13	16	19	19	22	16	17
80% to < 95%	1	4	3	4	8	4	4	5
40% to < 80%	2	2	2	1	2	3	1	2
< 40%	0	0	0	0	1	1	1	0
<b>Total (n=)</b>	<b>26</b>	<b>19</b>	<b>21</b>	<b>24</b>	<b>30</b>	<b>30</b>	<b>22</b>	<b>24</b>

Indicator 11: **Resistance Testing and Results**

Suppressed	1187	1197	1135	1163	1267	1235	1232	1208
Wild Type	189	178	195	173	156	147	152	152
Never Genotyped	7	8	8	9	6	3	8	9
1-Class	41	33	37	27	20	21	16	21
2-Class	7	4	8	3	4	3	3	3
3-Class	1	2	2	0	1	0	0	1
4-Class	0	0	0	0	0	0	0	0
<b>Total (n=)</b>	<b>1432</b>	<b>1422</b>	<b>1385</b>	<b>1375</b>	<b>1454</b>	<b>1409</b>	<b>1411</b>	<b>1394</b>

Indicator 12: **AIDS-Defining Illness**

	2008	2009	2010	2011	2012	2013	2014	2015
CD4 < 200 at Cases	51	31	33	33	25	19	21	25
ART initiation <i>Rate per 100,000</i>	3.3	2.0	2.0	2.0	1.5	1.1	1.2	1.4
AIDS Cases Cases	26	17	24	22	14	19	13	23
(DTP Reports) <i>Rate per 100,000</i>	1.7	1.1	1.5	1.3	0.8	1.1	0.8	1.3
AIDS Cases Cases	30	20	29	24	14	11	15	19
(BCCDC Reports) <i>Rate per 100,000</i>	1.9	1.3	1.8	1.5	0.8	0.7	0.9	1.1

Indicator 13: **HIV-Related Mortality**

	2004	2005	2006	2007	2008	2009	2010	2011
British Columbia	105	146	142	100	79	63	54	59
Per 100 HIV+ Population	1.03	1.40	1.34	0.93	0.72	0.56	0.47	0.50
Per 100,000 Population	2.50	3.43	3.29	2.28	1.80	1.41	1.19	1.29